

## ADHD Self-Test

Is it possible that I have symptoms of ADHD?

Here is an ADHD test to give you an idea if you may have ADHD.

**Rate Yourself on a scale of 1 to 5, 5 being most often.**

**Six or more symptoms in each category must be present with a score of 3 or more.**

### **Inattention (low attention span)**

Six or more of the following symptoms of inattention have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level:

- Often fails to give close attention to details or makes careless mistakes in schoolwork or other activities.
- Often has difficulty sustaining attention in tasks or play activities.
- Often does not seem to listen when spoken to directly
- Often does not follow through on instruction and fails to finish schoolwork, chores or duties in the workplace (not due to oppositional behavior or failure to understand instructions.)
- Often has difficulty organizing tasks and activities.
- Often avoids, dislikes or is reluctant to engage in tasks that require sustained mental effort (such as schoolwork or homework.)
- Often loses things necessary for tasks or activities (e.g., toys school assignments, pencils, books, or tools.)
- Is often easily distracted by extraneous stimuli.
- Is often forgetful in daily activities.

### **Hyperactivity-Impulsiveness**

Six or more of the following symptoms of hyperactivity-impulsivity have persisted for at least 6 months to a degree that is maladaptive and inconsistent with developmental level:

- Often fidgets with hands or feet or squirms in seat.
- Often leaves seat in classroom or in other situation in which remaining seated is expected.
- Often runs about or climbs excessively in situations in which it is inappropriate (in adolescents or adults may be limited to subjective feelings of restlessness.)
- Often has difficulty playing or engaging in leisure activities quietly.
- Is often "on the go" or often acts as if "driven by a motor."
- Often talks excessively.
- Often blurts out answers before questions have been completed.
- Often has difficulty awaiting turn.
- Often interrupts or intrudes on others (e.g., at school or work and at home.)

### **ADHD Test Additional Considerations**

- Some hyperactive-impulsive and inattentive symptoms that caused impairment were present before age 7 years.
- Some impairment from the symptoms is present in two or more settings (e.g., at school or work and at home.)
- There must be clear evidence of clinically significant impairment in social, academic or occupational functioning.
- The symptoms do not occur exclusively during the course of a Pervasive Developmental Disorder, Schizophrenic or other Psychotic Disorder and are not better accounted for by another mental disorder (e.g., Mood Disorder, Anxiety Disorder, Dissociate Disorder or a Personality Disorder.)

## General Adult ADHD Test Symptom Checklist

In conjunction with other diagnostic techniques the following general adult ADHD checklist helps further define ADHD symptoms. No ADHD adult has all of the symptoms, but if you notice a strong presence of more than 20 of these symptoms, there is a strong likelihood of ADHD.

Please read this list of behaviors and rate yourself (or the person who has asked you to rate him or her) on each behavior listed. Print the text below and then, using the following scale, and place the appropriate number next to the item. The scoring procedure is at the end of the test.

**0 = Never    1 = Rarely    2 = Occasionally    3 = Frequently    4 = Very Frequently**

**IMPORTANT:** This is not a tool for self-diagnosis. Its purpose is simply to help you determine whether ADHD may be a factor in the behavior of the person you are assessing using this checklist. An actual diagnosis can be made only by an experienced professional.

### Past History

1. \_\_\_\_\_ History of ADHD symptoms in childhood, such as distractibility, short attention span, impulsiveness or restlessness. ADHD doesn't start at age 30.
2. \_\_\_\_\_ History of not living up to potential in school or work (report cards with comments such as not living up to potential)
3. \_\_\_\_\_ History of frequent behavior problems in school (mostly for males)
4. \_\_\_\_\_ History of bed-wetting past age 5
5. \_\_\_\_\_ Family history of ADD, learning problems, mood disorders or substance abuse problems

### Short Attention Span/Distractibility

6. \_\_\_\_\_ Short attention span, unless very interested in something
7. \_\_\_\_\_ Easily distracted, tendency to drift away (although at times can be hyper focused)
8. \_\_\_\_\_ Lacks attention to detail, due to distractibility
9. \_\_\_\_\_ Trouble listening carefully to directions
10. \_\_\_\_\_ Frequently misplaces things
11. \_\_\_\_\_ Skips around while reading or goes to the end first, trouble staying on track
12. \_\_\_\_\_ Difficulty learning new games because it is hard to stay on track during directions
13. \_\_\_\_\_ Easily distracted during sex causing frequent breaks or turnoffs during love making
14. \_\_\_\_\_ Poor listening skills
15. \_\_\_\_\_ Tendency to be easily bored (tunes out)

### Restlessness

16. \_\_\_\_\_ Restlessness, constant motion, legs moving, fidgety
17. \_\_\_\_\_ Has to be moving in order to think
18. \_\_\_\_\_ Trouble sitting still, such as trouble sitting in one place for too long, sitting at a desk job for long periods, sitting through a movie
19. \_\_\_\_\_ An internal sense of anxiety or nervousness

### Impulsiveness

20. \_\_\_\_\_ Impulsive in words and/or actions (spending)
21. \_\_\_\_\_ Say just what comes to mind without considering its impact (tactless)
22. \_\_\_\_\_ Trouble going through established channels, trouble following proper procedure, an attitude of, "Read the directions only if all else fails."
23. \_\_\_\_\_ Impatient, low frustration tolerance
24. \_\_\_\_\_ A prisoner of the moment
25. \_\_\_\_\_ Frequent traffic violations
26. \_\_\_\_\_ Frequent, impulsive job changes
27. \_\_\_\_\_ Tendency to embarrass others
28. \_\_\_\_\_ Lying or stealing on impulse

### **Poor Organization**

29. \_\_\_\_\_ Poor organization and planning, trouble maintaining an organized work/living area
30. \_\_\_\_\_ Chronically late or chronically in a hurry
31. \_\_\_\_\_ Often has piles of stuff
32. \_\_\_\_\_ Easily overwhelmed by tasks of daily living
33. \_\_\_\_\_ Poor financial management (late bills, checkbook a mess, spending unnecessary money on late fees)
34. \_\_\_\_\_ Some adults with ADHD are very successful, but often only if they are surrounded with people who organize them.

### **Problems Getting Started and Following Through**

35. \_\_\_\_\_ Chronic procrastination or trouble getting started
36. \_\_\_\_\_ Starting projects but not finishing them, poor follow through
37. \_\_\_\_\_ Enthusiastic beginnings but poor endings
38. \_\_\_\_\_ Spends excessive time at work because of inefficiencies
39. \_\_\_\_\_ Inconsistent work performance

### **Negative Internal Feelings**

40. \_\_\_\_\_ Chronic sense of underachievement, feeling you should be much further along in your life
41. \_\_\_\_\_ Chronic problems with self-esteem
42. \_\_\_\_\_ Sense of impending doom
43. \_\_\_\_\_ Mood swings
44. \_\_\_\_\_ Negativity
45. \_\_\_\_\_ Frequent feeling of demoralization or that things won't work out for you

### **Relational Difficulties**

46. \_\_\_\_\_ Trouble sustaining friendships or intimate relationships, promiscuity
47. \_\_\_\_\_ Trouble with intimacy
48. \_\_\_\_\_ Tendency to be immature
49. \_\_\_\_\_ Self-centered; immature interests
50. \_\_\_\_\_ Failure to see others' needs or activities as important
51. \_\_\_\_\_ Lack of talking in a relationship
52. \_\_\_\_\_ Verbally abusive to others
53. \_\_\_\_\_ Proneness to hysterical outburst
54. \_\_\_\_\_ Avoids group activities
55. \_\_\_\_\_ Trouble with authority

### **Short Fuse**

56. \_\_\_\_\_ Quick responses to slights that are real or imagined
57. \_\_\_\_\_ Rage outbursts, short fuse

### **Frequent Search For High Stimulation**

58. \_\_\_\_\_ Frequent search for high stimulation (bungee jumping, gambling, high stress jobs, ER doctors, doing many things at once, etc.)
59. \_\_\_\_\_ Tendency to seek conflict, be argumentative or to start disagreements for the fun of it

### **Tendency To Get Stuck (thoughts or behaviors)**

60. \_\_\_\_\_ Tendency to worry needlessly and endlessly
61. \_\_\_\_\_ Tendency toward addictions (food, alcohol, drugs, work)

**Switches Things Around**

- 62. \_\_\_\_\_ Switches around numbers, letters or words
- 63. \_\_\_\_\_ Turn words around in conversations

**Writing/Fine Motor Coordination Difficulties**

- 64. \_\_\_\_\_ Poor writing skills (hard to get information from brain to pen)
- 65. \_\_\_\_\_ Poor handwriting, often prints
- 66. \_\_\_\_\_ Coordination difficulties

**The Harder I Try The Worse It Gets**

- 67. \_\_\_\_\_ Performance becomes worse under pressure.
- 68. \_\_\_\_\_ Test anxiety or during tests your mind tends to go blank
- 69. \_\_\_\_\_ The harder you try, the worse it gets
- 70. \_\_\_\_\_ Work or schoolwork deteriorates under pressure
- 71. \_\_\_\_\_ Tendency to turn off or become stuck when asked questions in social situations
- 72. \_\_\_\_\_ Falls asleep or becomes tired while reading

**Sleep/Wake Difficulties**

- 73. \_\_\_\_\_ Difficulties falling asleep, may be due to too many thoughts at night
- 74. \_\_\_\_\_ Difficulty coming awake (may need coffee or other stimulant or activity before feeling fully awake.)

**Low Energy**

- 75. \_\_\_\_\_ Periods of low energy, especially early in the morning and in the afternoon
- 76. \_\_\_\_\_ Frequently feeling tired

**Sensitive To Noise Or Touch**

- 77. \_\_\_\_\_ Easily startled
- 78. \_\_\_\_\_ Sensitive to touch, clothing, noise and light

---

**When you have completed the above ADHD test, calculate the following:**

- 1. Total Score: \_\_\_\_\_
- 2. Total Number of Items with a score of three (3) or more: \_\_\_\_\_
- 3. Score for Item #1: \_\_\_\_\_
- 4. Score for Item #6: \_\_\_\_\_
- 5. Score for Item #7: \_\_\_\_\_

More than 20 items with a score of three or more indicates a strong tendency toward ADHD. Items 1, 6, and 7 are essential to make the diagnosis.

Although this, or no self-test, is conclusive, if you need help you should consider contacting an iTherapyRX counselor or another health care professional.